

Amendments to the Claims

Please cancel Claims 5-7, 13, 19-49. Please amend Claims 1, 8 and 14. The Claim Listing below will replace all prior versions of the claims in the application:

Claim Listing

1. (Currently amended) A method for identifying an FGF receptor ligand comprising:
 - a) providing a cDNA expression library from an organism of interest;
 - b) providing a population of cells that express a DNA sequence encoding a heterologous FGF receptor that comprises an extracellular domain, a transmembrane domain, and an intracellular domain that is characterized by protein tyrosine kinase activity;
 - c) transforming the population of cells with the cDNA expression library; and
 - d) detecting protein kinase activity in clonally-derived cells, wherein elevated FGF receptor tyrosine kinase activity indicates the presence of an FGF receptor ligand.
2. (Original) The method according to Claim 1, wherein the cell is a yeast cell.
3. (Original) The method according to Claim 1, wherein the DNA sequence encoding a heterologous FGF receptor is carried on a CEN-based plasmid.
4. (Original) The method according to Claim 1, wherein the DNA sequence encoding a heterologous FGF receptor is inserted into a chromosome.
- 5-7. (Cancel)
8. (Currently amended) A method for identifying an FGF receptor ligand comprising:
 - a) providing a cDNA expression library from an organism of interest under the control of an inducible promoter;

- b) providing a population of cells that express a DNA sequence encoding a heterologous FGF receptor that comprises an extracellular domain, a transmembrane domain, and an intracellular domain that is characterized by protein tyrosine kinase activity;
 - c) transforming the population of cells with the cDNA expression library; and
 - d) detecting protein kinase activity in clonally-derived cells, wherein elevated FGF receptor tyrosine kinase activity indicates the presence of an FGF receptor ligand.
9. (Original) The method according to Claim 8, wherein the cell is a yeast cell.
10. (Original) The method according to Claim 8, wherein the DNA sequence encoding a heterologous FGF receptor is carried on a CEN-based plasmid.
11. (Original) The method according to Claim 8, wherein the DNA sequence encoding a heterologous FGF receptor is inserted into a chromosome.
12. (Original) The method according to Claim 8, wherein the DNA sequence encoding a heterologous FGF receptor is constitutively expressed.
13. (Cancel)
14. (Currently amended) A method for identifying an FGF receptor ligand comprising:
- a) providing a cDNA expression library from an organism of interest;
 - b) providing a population of cells that constitutively express a DNA sequence encoding a heterologous FGF receptor that comprises an extracellular domain, a transmembrane domain, and an intracellular domain that is characterized by protein tyrosine kinase activity;
 - c) transforming the population of cells with the cDNA expression library; and
 - d) detecting protein kinase activity in clonally-derived cells, wherein elevated FGF receptor tyrosine kinase activity indicates the presence of an FGF receptor ligand.

15. (Original) The method according to Claim 14, wherein the cell is a yeast cell.
16. (Original) The method according to Claim 14, wherein the DNA sequence encoding a heterologous FGF receptor is carried on a CEN-based plasmid.
17. (Original) The method according to Claim 14, wherein the DNA sequence encoding a heterologous FGF receptor is inserted into a chromosome.
18. (Original) The method according to Claim 14, wherein the cDNA expression library having polynucleotide inserts under the control of an inducible promoter.
- 19-49. (Cancel)